

Non-Abrikosov Vortex and Topological Knot in Two-gap Superconductor

Y. M. Cho* and Pengming Zhang†

Center for Theoretical Physics and School of Physics
College of Natural Sciences,
Seoul National University, Seoul 151-742, Korea

We establish the existence of topologically stable knot in two-gap superconductor whose topology $\pi_3(S^2)$ is fixed by the Chern-Simon index of the electromagnetic potential. We present a helical magnetic vortex solution in Ginzburg-Landau theory of two-gap superconductor which has a non-vanishing condensate at the core, and identify the knot as a twisted magnetic vortex ring made of the helical vortex. We discuss how the knot can be constructed in the recent two-gap MgB₂ superconductor.

PACS numbers: 74.20.-z, 74.20.De, 74.60.Ge, 74.60.Jg, 74.90.+n

Keywords: twisted magnetic vortex ring, topological knot in two-gap superconductor, Chern-Simon topology of magnetic field

Topological objects, in particular finite energy topological objects have played important role in physics. In condensed matter physics the best known topological object is the Abrikosov vortex in one-gap superconductors (and similar ones in Bose-Einstein condensates and superfluids), which have been the subject of intensive studies. But the advent of two-component BEC and two-gap superconductors [1, 2] has opened a new possibility for us to have far more interesting topological objects in condensed matters [3, 4, 5, 6, 7]. *The purpose of this Letter is two-fold; to present a new type of non-Abrikosov vortex which has a non-vanishing condensate at the core and to establish the existence of a stable knot in Abelian two-gap superconductors.* We first construct a new type of non-Abrikosov vortex which has a non-vanishing condensate at the core in Ginzburg-Landau theory of two-gap superconductor, and show that it can be twisted to form a helical magnetic vortex. With this we show that the twisted vortex ring made of the helical magnetic vortex becomes a topological knot whose knot topology $\pi_3(S^2)$ is fixed by the Chern-Simon index of the electromagnetic potential. The knot has two supercurrents, the electromagnetic current around the knot tube which confines the magnetic flux and the neutral current along the knot which generates a net angular momentum which prevents the collapse of the knot, This means that the knot has a dynamical (as well as topological) stability.

There have been two objections against the existence of a stable magnetic vortex ring in Abelian superconductors. First, it is supposed to be unstable due to the tension created by the ring [8]. Indeed if one constructs a vortex ring from an Abrikosov vortex, it becomes unstable because of the tension. On the other hand, if we first twist the magnetic vortex making it periodic in z -

coordinate and then make a vortex ring connecting the periodic ends together, it should become a stable vortex ring. This is because the non-trivial twist of the magnetic field forbids the untwisting of the vortex ring by any smooth deformation of field configuration. The other objection is that the Abelian gauge theory is supposed to have no non-trivial knot topology which allows a stable vortex ring. This again is a common misconception. Actually the theory has a well-defined knot topology $\pi_3(S^2)$ described by the Chern-Simon index of the electromagnetic potential. So *a priori* there is no reason whatsoever why the Abelian superconductor can not have a topological knot. The real question then is whether the dynamics of the superconductor can actually allow such a knot configuration. In the following we show that indeed a two-gap Abelian superconductor does.

In mean field approximation the Hamiltonian of the Ginzburg-Landau theory of two-gap superconductor could be expressed by [7]

$$\mathcal{H} = \frac{\hbar^2}{2m_1} |(\nabla + ig\mathbf{A})\tilde{\phi}_1|^2 + \frac{\hbar^2}{2m_2} |(\nabla + ig\mathbf{A})\tilde{\phi}_2|^2 + \tilde{V}(\tilde{\phi}_1, \tilde{\phi}_2) + \frac{1}{2}(\nabla \times \mathbf{A})^2, \quad (1)$$

where $\tilde{V}$ is the effective potential. One can simplify the above Hamiltonian with a proper normalization of $\tilde{\phi}_1$ and $\tilde{\phi}_2$ to $\phi_1 = (\hbar/\sqrt{2m_1})\tilde{\phi}_1$ and $(\hbar/\sqrt{2m_2})\tilde{\phi}_2$,

$$\mathcal{H} = |(\nabla + ig\mathbf{A})\phi|^2 + V(\phi) + \frac{1}{2}(\nabla \times \mathbf{A})^2, \quad (2)$$

where $\phi = (\phi_1, \phi_2)$ and V is the normalized potential. A most general quartic potential which has the $U(1) \times U(1)$ symmetry is given by

$$V = \frac{\lambda_{11}}{2} |\phi_1|^4 + \lambda_{12} |\phi_1|^2 |\phi_2|^2 + \frac{\lambda_{22}}{2} |\phi_2|^4 - \mu_1 |\phi_1|^2 - \mu_2 |\phi_2|^2, \quad (3)$$

*Electronic address: ymcho@yongmin.snu.ac.kr

†Electronic address: zhpm@phya.snu.ac.kr

where λ_{ij} are the quartic coupling constants and μ_i are the chemical potentials. But in this paper we will adopt the following potential for simplicity

$$V = \frac{\lambda}{2} \left((\phi^\dagger \phi) - \frac{\mu^2}{\lambda} \right)^2 + \alpha |\phi_1|^2 |\phi_2|^2. \quad (4)$$

Notice that when $\alpha = 0$, the Hamiltonian (2) has a global $SU(2)$ symmetry as well as the local $U(1)$ symmetry, so that the α -term can be viewed as a symmetry breaking term of $SU(2)$ to $U(1)$. We choose (4) simply because the existence of the topological objects we discuss in this paper is not so sensitive to the detailed form of the potential.

With the above Hamiltonian one may try to obtain a non-Abelian vortex. With

$$\begin{aligned} \phi &= \frac{1}{\sqrt{2}} \rho \xi, \quad \xi^\dagger \xi = 1, \quad \hat{n} = \xi^\dagger \vec{\sigma} \xi, \\ C_\mu &= \frac{2i}{g} \xi^\dagger \partial_\mu \xi, \quad \rho_0 = \frac{2\mu^2}{\lambda}, \end{aligned} \quad (5)$$

we have the following equations of motion [3]

$$\begin{aligned} \partial^2 \rho - \left(\frac{1}{4} (\partial_\mu \hat{n})^2 + g^2 (A_\mu - \frac{1}{2} C_\mu)^2 \right) \rho &= \frac{\lambda}{2} (\rho^2 - \rho_0^2) \rho + \alpha \rho^3 \xi_1^* \xi_1 \xi_2^* \xi_2, \\ \hat{n} \times \partial^2 \hat{n} + 2 \frac{\partial_\mu \rho}{\rho} \hat{n} \times \partial_\mu \hat{n} - \frac{2}{g \rho^2} \partial_\mu F_{\mu\nu} \partial_\nu \hat{n} &= \frac{\alpha}{2} (\hat{k} \cdot \hat{n}) \rho^2 \hat{k} \times \hat{n}, \\ \partial_\mu F_{\mu\nu} &= g^2 \rho^2 (A_\mu - \frac{1}{2} C_\mu), \end{aligned} \quad (6)$$

where $\vec{\sigma}$ is the Pauli matrix and $\hat{k} = (0, 0, 1)$. Notice that it has the vacuum at $\rho = \rho_0$, and $\xi = (1, 0)$ or $\xi = (0, 1)$.

The equation allows two conserved currents, the electromagnetic current j_μ and the neutral current k_μ ,

$$\begin{aligned} j_\mu &= g^2 \rho^2 (A_\mu - \frac{1}{2} C_\mu) \\ k_\mu &= g^2 \rho^2 \left[A_\mu (\xi_1^* \xi_1 - \xi_2^* \xi_2) + \frac{i}{g} (\partial_\mu \xi_1^* \xi_1 \right. \\ &\quad \left. + \xi_2^* \partial_\mu \xi_2) \right], \end{aligned} \quad (7)$$

which are nothing but the Noether currents of the $U(1) \times U(1)$ symmetry of the Hamiltonian (2). Indeed they are the sum and difference of two electromagnetic currents of ϕ_1 and ϕ_2

$$j_\mu = j_\mu^{(1)} + j_\mu^{(2)}, \quad k_\mu = j_\mu^{(1)} - j_\mu^{(2)}. \quad (8)$$

Clearly the conservation of j_μ follows from the last equation of (6). But the conservation of k_μ comes from the second equation of (6), which (together with the last equation) tells the existence of a partially conserved $SU(2)$ current $\vec{j}_\mu$,

$$\vec{j}_\mu = g \rho^2 \left(\frac{1}{2} \vec{n} \times \partial_\mu \vec{n} - g (A_\mu - \frac{1}{2} C_\mu) \vec{n} \right). \quad (9)$$

This $SU(2)$ current is exactly conserved when $\alpha = 0$. But notice that $k_\mu = \hat{k} \cdot \vec{j}_\mu$. This assures the conservation of k_μ even when $\alpha \neq 0$. It is interesting to notice that j_μ and k_μ are precisely the $\hat{n}$ and $\hat{k}$ components of $\vec{j}_\mu$.

To obtain the desired knot we construct a helical magnetic vortex first [3]. Let (ϱ, φ, z) be the cylindrical coordinates and let

$$\begin{aligned} \rho &= \rho(\varrho), \quad \xi = \begin{pmatrix} \cos \frac{f(\varrho)}{2} \exp(-in\varphi) \\ \sin \frac{f(\varrho)}{2} \exp(ikz) \end{pmatrix}, \\ A_\mu &= \frac{1}{g} \left(n A_1(\varrho) \partial_\mu \varphi + m k A_2(\varrho) \partial_\mu z \right), \end{aligned} \quad (10)$$

and find

$$\begin{aligned} \hat{n} &= \begin{pmatrix} \sin f(\varrho) \cos(n\varphi + mkz) \\ \sin f(\varrho) \sin(n\varphi + mkz) \\ \cos f(\varrho) \end{pmatrix}, \\ C_\mu &= \frac{\cos f(\varrho) + 1}{g} n \partial_\mu \varphi + \frac{\cos f(\varrho) - 1}{g} m k \partial_\mu z, \\ j_\mu &= g \rho^2 \left(n \left(A_1 - \frac{\cos f + 1}{2} \right) \partial_\mu \varphi \right. \\ &\quad \left. + m k \left(A_2 - \frac{\cos f - 1}{2} \right) \partial_\mu z \right), \\ k_\mu &= g \rho^2 \left(n \left(A_1 \cos f - \frac{\cos f + 1}{2} \right) \partial_\mu \varphi \right. \\ &\quad \left. + m k \left(A_2 \cos f + \frac{\cos f - 1}{2} \right) \partial_\mu z \right). \end{aligned} \quad (11)$$

With this (6) becomes

$$\begin{aligned} \ddot{\rho} + \frac{1}{\varrho} \dot{\rho} - \left[\frac{1}{4} \left(\dot{f}^2 + \left(\frac{n^2}{\varrho^2} + m^2 k^2 \right) \sin^2 f \right) \right. \\ \left. + \frac{n^2}{\varrho^2} \left(A_1 - \frac{\cos f + 1}{2} \right)^2 + m^2 k^2 \left(A_2 - \frac{\cos f + 1}{2} \right)^2 \right] \rho \\ = \frac{\lambda}{2} (\rho^2 - \rho_0^2) \rho + \frac{\alpha}{4} \rho^3 \sin^2 f, \\ \ddot{f} + \left(\frac{1}{\varrho} + 2 \frac{\dot{\rho}}{\rho} \right) \dot{f} - 2 \left(\frac{n^2}{\varrho^2} (A_1 - \frac{1}{2}) \right. \\ \left. + m^2 k^2 (A_2 + \frac{1}{2}) \right) \sin f = \frac{\alpha}{4} \rho^2 \sin 2f, \\ \ddot{A}_1 - \frac{1}{\varrho} \dot{A}_1 - g^2 \rho^2 \left(A_1 - \frac{\cos f + 1}{2} \right) = 0, \\ \ddot{A}_2 + \frac{1}{\varrho} \dot{A}_2 - g^2 \rho^2 \left(A_2 - \frac{\cos f - 1}{2} \right) = 0. \end{aligned} \quad (12)$$

Now, we impose the following boundary condition for the non-Abelian vortex,

$$\begin{aligned} \dot{\rho}(0) &= 0, \quad \rho(\infty) = \rho_0, \quad f(0) = \pi, \quad f(\infty) = 0, \\ A_1(0) &= 0, \quad A_1(\infty) = 1, \\ A_2(0) &= 0, \quad A_2(\infty) = 0. \end{aligned} \quad (13)$$

This need some explanation. First, notice that here we have $\dot{\rho}(0) = 0$, not $\rho(0) = 0$. This would have been unacceptable in one-gap superconductor because this would




FIG. 1: The helical vortex (solid line) with $m = n = 1$ and the straight vortex (dotted line) $m = 0$ and $n = 1$ in two-gap superconductor. Here we have put $g = 1$, $\lambda = 2$, $k = 0.1 \rho_0$, $\alpha = 0.01$, and ϱ is in the unit of $1/\rho_0$. Notice that the straight vortex has no A_2 .

create a singularity at the core of the vortex. But with the ansatz (10) the boundary condition allows a smooth vortex in two-gap superconductor. Secondly, we have $A_2(0) = 0$ instead of a condition on $A_2(0)$. It turns out that this is the correct way to obtain a smooth solution. Indeed we find that $A_2(0)$ is determined by k .

With the boundary condition we can integrate (12) and obtain the non-Abelian vortex solution of the two-gap superconductor which is shown in Fig.1. Notice that we have plotted $\rho_1 = \rho \cos f/2$ and $\rho_2 = \rho \sin f/2$ which represent the density of ϕ_1 and ϕ_2 , instead of ρ and f , in the figure. There are three points that should be emphasized here. First, the doublet ϕ starts from the second component at the core, but the first component takes over completely at the infinity. This is due to the boundary condition $f(0) = \pi$ and $f(\infty) = 0$, which assures that our solution describes a new type of non-Abriksov vortex which has a non-vanishing condensate at the core [3]. Secondly the electromagnetic current j_μ of the vortex is helical, so that it creates a helical magnetic flux which has a quantized flux $\Phi_z = 2\pi n/g$ around the z -axis as well as a (fractionally) quantized flux $\Phi_\phi = -2\pi m A_2(0)/g$ along the z -axis. And obviously the two magnetic fluxes are linked together. But it creates a net current i_ϕ only around the z -axis, with $i_z = i_z^{(1)} + i_z^{(2)} = 0$. Finally the vortex has another (neutral) helical k_μ , which has a net current k_z along the z -axis (as well as k_ϕ around the vortex). Notice that $k_z = i_z^{(1)} - i_z^{(2)} = 2i_z^{(1)}$. This is because the two electromagnetic currents $i_z^{(1)}$ and $i_z^{(2)}$ of ϕ_1 and ϕ_2 are equal but flow oppositely, which generates a non-vanishing k_z . This is why we call k_μ a neutral current. In this sense the vortex can be viewed as superconducting, even though the net electromagnetic current along the knot is zero. The two helical supercurrents j_μ and k_μ are




FIG. 2: The helical supercurrents with $m = n = 1$ in two-gap superconductor. Here we have put $g = 1$, $\lambda = 2$, $k = 0.1 \rho_0$, $\alpha = 0.01$, and ϱ is in the unit of $1/\rho_0$.

plotted in Fig. 2.

Of course, the helical magnetic vortex has only a heuristic value, because it will be untwisted unless the periodicity condition is enforced by hand. But if we make a vortex ring by smoothly bending and connecting two periodic ends, the periodicity condition is naturally enforced. And the twisted magnetic vortex ring becomes a topological knot [3]. There are two ways to describe the knot topology. First, the doublet ξ (in the vortex ring) acquires a non-trivial topology $\pi_3(S^2)$ which provides the knot quantum number,

$$Q = -\frac{1}{4\pi^2} \int \epsilon_{ijk} \xi^\dagger \partial_i \xi (\partial_j \xi^\dagger \partial_k \xi) d^3x \\ = \frac{g^2}{32\pi^2} \int \epsilon_{ijk} C_i (\partial_j C_k - \partial_k C_j) d^3x = mn. \quad (14)$$

This is nothing but the Chern-Simon index of the potential C_μ . Perhaps more importantly, the knot topology can also be described by the Chern-Simon index of the electromagnetic potential A_μ . Indeed we have

$$Q = \frac{g^2}{8\pi^2} \int \epsilon_{ijk} A_i F_{jk} d^3x = mn, \quad (15)$$

which describes the twisted knot topology of the real (physical) magnetic flux. It describes the linking number of two quantized magnetic fluxes Φ_z and Φ_ϕ . Obviously two flux rings linked together can not be separated by any continuous deformation of the field configuration. This provides the topological stability of the knot.

Furthermore the topological stability is backed up by a dynamical stability. This is because the knot is made of two quantized magnetic fluxes linked together. And the flux trapped inside of the knot ring can not be squeezed out, and provides a repulsive force against the collapse of the knot. This makes the knot dynamically stable. An-




FIG. 3: (Color online). The energy profile of superconducting knot with $m = n = 1$. Here we have put $g = 1$, $\lambda = 2$, $k = 0.1 \rho_0$, $\alpha = 0.01$.

other way to understand this dynamical stability is to notice that the neutral supercurrent k_z of the helical vortex becomes a supercurrent along the knot, which generates a net angular momentum around the knot. And this provides the centrifugal repulsive force preventing the knot to collapse. This makes the knot dynamically stable. We can find the energy profile of the twisted magnetic vortex ring numerically. Minimizing the energy we obtain the energy profile of the knot shown in Fig. 3.

Clearly the dynamical stability of the knot comes from the helical structure of the vortex. But this helical structure is precisely what provides the knot topology to the vortex ring. The nontrivial topology expresses itself in the form of the helical magnetic flux which provides the dynamical stability of the knot. Conversely the helical magnetic flux assures the existence of the knot topology which guarantees the topological stability of the knot. It is this remarkable interplay between dynamics and topology which assures the existence of the stable superconducting knot in two-gap superconductor.

We close with the following remarks:

1. The existence of a knot in two-gap superconductor has been conjectured before [3, 7]. In fact, similar knots have been asserted to exist almost everywhere in physics, in atomic physics in two-component Bose-Einstein con-

densates [3, 5, 6], in nuclear physics in Skyrme theory [9, 10], in high energy physics in QCD [11]. In this paper we have provided a concrete evidence for the existence of a stable knot in two-gap superconductor, which can be interpreted as a twisted magnetic vortex ring.

2. In this Letter we have concentrated on the Abelian gauge theory of two-gap superconductor, which is made of two condensates of same charge. But we emphasize that a similar knot should also exist in the non-Abelian gauge theory of two-gap superconductor which describes two condensates of opposite charge [4]. This is because two theories are mathematically identical to each other. This implies that such a knot can also exist in the liquid metallic hydrogen (LMH), because it can also be viewed as a non-Abelian two-gap superconductor made of opposite charge (the proton pairs and the electron pairs) [12].

3. One should try to construct the topological knot experimentally in two-gap superconductors. But before one does that, one must first confirm the existence of the non-Abrikosov vortex which has a non-vanishing condensate (the second component) at the core. This is very important because this will prove the existence of a new type of magnetic vortex in two-gap superconductor. With this confirmation one may try to construct the helical vortex and the knot. With some experimental ingenuity one should be able to construct these topological objects in MgB_2 or LMH.

As we have remarked there is a wide class of realistic potentials which allows similar non-Abrikosov vortex and topological knot. The detailed discussion on the subject and its applications in the presence of a more general potential which includes the Josephson interaction will be discussed separately [13].

ACKNOWLEDGEMENT

The work is supported in part by the ABRL Program of Korea Science and Engineering Foundation (Grant R14-2003-012-01002-0), and by the BK21 Project of the Ministry of Education.

-
- [1] C. Myatt *et al.*, Phys. Rev. Lett. **78**, 586 (1997); D. Stamper-Kurn, *et al.*, Phys. Rev. Lett. **80**, 2027 (1998).
 - [2] J. Nagamatsu *et al.*, Nature **410**, 63 (2001); S. L. Bud'ko *et al.*, Phys. Rev. Lett. **86**, 1877 (2001); C. U. Jung *et al.*, Appl. Phys. Lett. **78**, 4157 (2001).
 - [3] Y. M. Cho, cond-mat/012325; Int. J. Pure Appl. Phys. **1**, 246 (2005).
 - [4] Y. M. Cho, cond-mat/012492; Phys. Rev. **B72**, 212516 (2005).
 - [5] U. Al Khawaja and H. Stoof, Nature **411**, 918 (2001); C. Savage and J. Ruostekoski, Phys. Rev. Lett. **91**, 010403 (2003).
 - [6] Y. M. Cho, H. J. Khim, and Pengming Zhang, Phys. Rev. **A72**, 063603 (2005).
 - [7] E. Babaev, Phys. Rev. Lett. **89**, 067001 (2002); E. Babaev, L. Faddeev, and A. Niemi, Phys. Rev. **B65**, 100512 (2002).
 - [8] K. Huang and R. Tipton, Phys. Rev. **D23**, 3050 (1981).
 - [9] L. Faddeev and A. Niemi, Nature **387**, 58 (1997); R. Battye and P. Sutcliffe, Phys. Rev. Lett. **81**, 4798 (1998).
 - [10] Y. M. Cho, Phys. Rev. Lett. **87**, 252001 (2001); Phys. Lett. **B603**, 88 (2004).
 - [11] Y. M. Cho, Phys. Lett. **B616**, 101 (2005).
 - [12] N. Ashcroft, Phys. Rev. Lett. **92**, 187002 (2004); E. Babaev, A. Sudbe, and N. Ashcroft, Nature **431**, 666 (2004).
 - [13] Y. M. Cho, and Pengming Zhang, to be published.